



SURAT PERNYATAAN

Saya yang bertanda tangan dibawah ini :

N a m a : Dr. Ir. Hj. Nurmala, MM
NIP : 195608151988112001
NIDN : 0015085608
Pangkat / Gol : Pembina Utama Muda/ IV.c
Alamat Kantor : Jl. Ahmad Yani Pontianak78124 Kalimantan Barat
No. Telp : 089653718456
Surel : ibu.nurmala@gmail.com

Dengan ini menerangkan bahwa jurnal artikel saya dibawah ini:

“Effects of ameliorant on the growth of snakehead fish (*Channa striata*) juvenile reared in acid sulfate water medium located in tidal land.”

(www.bioflux.com.ro/docs/2018.1919-1926)

Memiliki persentasi tingkat plagiarisme yang tinggi (100%). Ini disebabkan ketika pengecekan persentasi plagiarisme dilakukan (menggunakan program Urkund setara Tunitin), artikel tersebut dalam bentuk Jurnal Daring (E Journal) yang bersifat open access dan pengecekan similarity dilakukan bukan dengan akun pribadi, melainkan dengan akun rekan satu lembaga sehingga terdeteksi sebagai artikel yang memiliki persentasi plagirisme yang tinggi. Terlampir hasil dan komentar Urkund terkait artikel tersebut.

Dan URL yang diberikan untuk chek similarity adalah repository institusi asal saya, yakni Politeknik Negeri Pontianak, untuk menjaga keamanan kerahasiaan account saya dalam mengakses program Urkund berbayar ini.

Demikian Surat Pernyataan ini dibuat untuk digunakan sebagaimana mestinya.

Pontianak, Pontianak, 12 Maret 2019

Yang menyatakan,

Dr. Ir. Hj. Nurmala, MM
NIP 195608151988112001

Urkund Analysis Result

Analysed Document: Artikel 2.pdf (D48356647)
Submitted: 2/26/2019 3:25:00 PM
Submitted By: n.simbolon@graduate.curtin.edu.au
Significance: 100 %

Sources included in the report:

<http://www.bioflux.com.ro/docs/2018.1919-1926.pdf>

Instances where selected sources appear:

1

0: <http://www.bioflux.com.ro/docs/2018.1919-1926.pdf>

100%

AAFL Bioflux, 2018, Volume 11, Issue 6. 1919 <http://www.bioflux.com.ro/aafl>

Effects of ameliorant on the growth of snakehead fish (*Channa striata*) juvenile reared in acid sulfate water

medium located in tidal land 1 Purnamawati, 1 Nurmala, 1 Mochammad I. Shilman, 2 Eko Dewantoro, 3 Diah A. S. Utami 1

Study Program of Aquaculture, Department of Marine Science and Fisheries, Pontianak State Polytechnic, Pontianak, West Kalimantan, Indonesia; 2 Study Program of Aquaculture, Faculty of Fisheries and Marine Science,

Muhammadiyah University of Pontianak, Pontianak, West Kalimantan, Indonesia; 3 Study Program of Aquaculture, Marine and Fisheries Polytechnic of Jembrana, Jembrana, Bali,

Indonesia. Corresponding author: Purnamawati, pur_polnep@yahoo.com Abstract. The aim of this study was to evaluate the effects of the use of

an ameliorant produced from banana stems to increase water pH and improve the growth of the snakehead fish (*Channa striata*). The experiment applied

a completely randomized design. The treatments applied in this study consisted of three treatments

including ameliorant (300 kg ha^{-1}), lime (CaCO_3) (180 kg ha^{-1}), and control (without ameliorant and lime). Each treatment applied four replicates.

C. striata juveniles with an average initial length of 2.5 ± 0.2 cm and an average initial weight of 0.27 ± 0.03 g were reared in 12 units of hapa net cages measuring $2 \times 3 \times 1.5$ m with a stocking density of 2 fish per L, for 60 days. The fish were fed commercial feed with a

protein content of 40%,

twice a day (morning and afternoon) until to apparent satiation. The

use of an ameliorant in the experimental medium significantly

gave better

results shown by higher survival (70.55%), specific growth rate ($6.42\% \text{ day}^{-1}$), feed efficiency (89.07%), protein retention (53.41%), fat retention (89.41%),

and albumin content ($3.12 \text{ g per } 100 \text{ mL}$) than those of control. Key Words: air-breathing fish, banana stems, dolomite, tidal area. Introduction. Indonesia has about 20.11 million ha of tidal swamp land consisting of 2.07 million ha of tidal potential land, 6.71 million ha of acid sulfate land, 10.89 million ha of peatland, and 0.44 million ha of saline land. The tidal land is

spread in several regions, including Sumatera, Kalimantan, Papua, and Sulawesi (Arsyad et al 2014). Tidal swamp land area of West Kalimantan is around 1.12 million ha, which is spread in several districts (BPS Kalbar 2012) and 20,532 ha of it is acid sulfate land (Alwi 2014).

Most of these lands have been used for fish farming, but in general, the productivity is still relatively low. The low productivity

of fisheries sectors in this area is due to various problems, especially poor water quality. In a low water quality condition, there are only certain fish species that can be cultivated, one of those species is the snakehead fish (*Channa striata*). *C. striata*, is a freshwater fish species, classified in Channidae family and Perciformes order (Nakkrasae et al 2015). This fish is a potential and important species to be developed as an aquaculture commodity (Mollah 1985; Marimuthu et al 2009; Mollah et al 2009; Rahman et al 2013), and has high economic value with prices ranging from 3.78 to 4.47 USD kg⁻¹. Moreover, this fish flesh is used as a treatment of post-surgical therapy and can increase body endurance (Gam et al 2006; Marimuthu et al 2009; Mohd & Abdul 2012). *C. striata* usually lives in rivers, rice fields and ponds. *C. striata* relatively prefers calm water (without agitation). Another advantage possessed by *C. striata* is its ability to

AACL Bioflux, 2018, Volume 11, Issue 6. 1920 <http://www.bioflux.com.ro/aac1>

live in low oxygen environment. But to grow optimally, this species needs to live in a medium with optimal water quality ranges. One of limiting factors that influences the *C. striata* cultivation in acid sulfate land is pH. Tidal land has a pH range of 2.53–3.39 (Purnamawati et al 2017a), sulfate concentration at 6.91–8.7 mg L⁻¹, dissolved oxygen less than 5 mg L⁻¹ (Purnamawati et al 2017b). This water quality condition can be improved by increasing the pH of the culture medium. To increase the water pH in fish ponds can be done by adding a certain amount of lime into fish ponds (de Queiroz et al 2004; Cavalcante et al 2012). However, the required lime is not always available, especially in some locations that are far from the centers of agricultural production facilities. To overcome this issue, some local based plant materials can be used as ameliorants to substitute the utilization of lime. Banana stem is a waste that can be used as an ameliorant. This is caused by banana stem has some advantages, including low cost, abundant availability, easy to be processed, and it is not harmful for the cultivated organisms. This material is expected to be a solution in aquaculture practices conducted in acid sulfate medium, especially in tidal land. This study aimed to evaluate the effectiveness of the use of banana stems as an ameliorant in acid sulfate medium in order to promote the growth of *C. striata* juveniles. Material and Method. The present

study was conducted on July-September 2018. The rearing of *C. striata* was conducted

in a tidal pond located in Kubu Raya District, West Kalimantan, Indonesia. The measurement of plasma glucose levels, albumin

levels, the proximate analysis of the fish feed and the experimental fish were

conducted

at Environmental and Food Technology Laboratory, Tanjungpura University, Pontianak, Indonesia. Experimental design. This study was conducted through a completely randomized design (CRD)

in a tidal pond using hapa net cages. The treatments applied in this study consisted of three treatments including ameliorant using banana stems (300 kg ha^{-1}), lime (CaCO_3) (180 kg ha^{-1}), and control (without ameliorant and lime).

Each treatment applied four replicates. Experimental fish. The experimental fish used in this study were *C. striata* juveniles.

The experimental fish had

an average initial length of $2.5 \pm 0.2 \text{ cm}$ and an average initial weight of $0.27 \pm 0.03 \text{ g}$.

Experimental mediums. The experimental mediums used in this study were 12 units of hapa net cage with a size of $2 \times 3 \times 1.5 \text{ m}$. The hapa net cages were set

in a pond, the top sides were covered with nets to avoid fish jumping out of the rearing cages. Experimental procedures.

The production of ameliorant from banana stems used a bio-activator (EM4: effective microorganism 4) to accelerate the composting process. The composting process occurred aerobically. After the raw material became a compost, it was then dried under sunlight. The chemical composition of a compost from banana stems consisted of Ca (1.54%), Mg (0.86%), C-organic (43.39%), N (2.48%), C/N (17.50), pH (8.8), KTK + ($41.07 \text{ cmol}^+ \text{ kg}^{-1}$), humic acid (2.50 ppm), fulvic acid (0.80 ppm), and moisture (68.78%) (Purnamawati 2017). Prior to the experiment, the experimental fish were acclimatized to experimental conditions. This step was carried out for 7 days in three net cages with a size of $4 \times 3 \times 1.5 \text{ m}$. *C. striata* were then reared in a tidal pond using hapa net cages with a stocking density of 2 fish per L (Vivekanandan 1977; Hidayatullah et al 2015). This study was conducted for 60 days. The experimental fish were fed commercial feed with a protein content of $\pm 40\%$, twice a day to apparent satiation.

Experimental parameters. The measurements of water quality parameters were performed following the procedures described by APHA (1989). The

water quality parameters measured were temperature, pH, dissolved oxygen, sulfate (SO_4^{2-}),

AAAL Bioflux, 2018, Volume 11, Issue 6. 1921 <http://www.bioflux.com.ro/aaal>

alkalinity, and

ammonia. The

measurements of temperature, pH and dissolved oxygen level were carried out every 15 days, while the measurements of sulfate, alkalinity and ammonia were performed at the beginning,

middle and the end of the experiment.

The measurement of albumin levels followed the method described by Infusino & Panteghini (2013), while the proximate analysis of the fish feed and the experimental fish were carried out according to the procedure by Takeuchi (1988). The measurement of plasma glucose levels was performed using a liquicolor glucose commercial kit GOD-PAP with calorimetric method and the results of the measurement were read with a spectrophotometer at a wavelength of 500 nm following the procedure

constructed by Wedemeyer & Yasutake (1977). The daily mortality was recorded to obtain the survival data.

Survival was calculated using a formula according to Kang'ombe & Brown (2008): $SR = (N_t \times N_0^{-1}) \times 100$ Note: SR = Survival (%) N_t = The number of the fish at the end of the study (individual) N_0 =

The number of the fish at the beginning of the

study (

individual) The fish weight samplings were done every 15 days to obtain the growth data (specific growth rate).

Specific growth rate was calculated using a formula stated by Weatherley & Gill (1989). $SGR = [(\ln W_2 - \ln W_1) / (t_2 - t_1)] \times 100$

Note:

SGR = Specific growth rate (% day⁻¹) W_1 = The average weight of the fish at the beginning of the study (g) W_2 = The average weight of the fish at time t_2 (g) $t_2 - t_1$ = Experimental duration Feed

intake (F) was obtained by summing the daily amount of the feed consumed during the study. The feed efficiency

was calculated using a formula constructed by Kang'ombe & Brown (2008). $e = [(W_t + D) - W_0] / F \times 100$

Note: e =

Feed efficiency W_0 = The weight of the fish at the beginning of the study (g) W_t = The weight of the fish at the end of the study (g) D = The weight of the dead fish during the study (g) F = The weight of total feed given during the study (g dry weight) Statistical analysis. The water

quality parameters were analyzed through descriptive statistic. Survival, specific growth rate, albumin content, feed efficiency, protein retention,

and fat retention were analyzed through ANOVA with a help of SPSS version 22

to see the differences among treatments. Results and Discussion. The

water quality ranges in the experimental mediums are shown in Table 1. Water quality ranges varied among one treatment and other treatments. Water temperature ranges in the experimental mediums were not relatively different among treatments and those were an optimal range for the fish growth. The pH, total alkalinity, and sulfate showed differences between ameliorant added and lime added treatments to control (without the addition of ameliorant and lime). This could be seen from the high pH and total alkalinity, as well as the low sulfate levels dissolved in water. There was no difference in dissolved oxygen (DO) and ammonia levels among ameliorant, lime and control treatments.

AACL

Bioflux, 2018, Volume 11, Issue 6. 1922 <http://www.bioflux.com.ro/aacl>

Table 1

The values of physical-chemical parameters of the water in each treatment during the study

Parameters	Treatments	Tolerance range *	and optimum range **
pH	Ameliorant	6.8–7.2	6.9–7.4
	Lime (Dolomite)	5.3–5.8	4.25–9.4*
	Control	4.25–9.4*	3)
Sulfate (SO ₄ ²⁻) (mg L ⁻¹)	Ameliorant	10.2–12.6	10.7–29.6
	Lime	20.1–48.2	5–150*
	Control	5–150*	1)
Alkalinity (mg L ⁻¹)	Ameliorant	85.1–115.8	71.0–110.4
	Lime	71.0–110.4	22.1–26.6
	Control	22.1–26.6	100–150*
Ammonia (NH ₃) (mg L ⁻¹)	Ameliorant	0.01–0.05	0.01–0.07
	Lime	0.01–0.07	0.03–0.05
	Control	0.03–0.05	0.54–1.57*
DO (mg L ⁻¹)	Ameliorant	4.12–5.80	4.15–5.47
	Lime	4.15–5.47	4.79–5.12
	Control	4.79–5.12	< 5
Temperature (°C)	Ameliorant	29.32–30.12	29.42–30.29
	Lime	29.42–30.29	28.32–30.02
	Control	28.32–30.02	26–32**

1) Wedemeyer (1996), 2) Qin et al (1997), 3) Courtenay, Jr. & Williams (2004), *) tolerance range, **) optimum range.

The growth performance (survival, SGR and feed efficiency) and physiological characteristics (protein retention, energy retention, and albumin content) of *C. striata* in 60-day-experiment can be seen in Table 2. The addition of an ameliorant into ponds located in an acid sulfate land could produce the highest specific growth rate (SGR) and albumin level of *C. striata* (6.42% day⁻¹ and 3.12 g per 100 mL, respectively), followed by lime added and control treatments ($p > 0.05$). The survival (SR) of *C. striata* juveniles ranged from 54.45 to 70.55%. The highest SR was obtained in ameliorant treatment, but it was not significantly different from the lime added treatment. The same results were also found in feed efficiency and protein retention (PR), while fat retention (FR) does not show a significant difference between among treatments tested in this study.

Table 2 Survival (SR), specific growth rate (SGR), albumin content, feed efficiency, protein retention (PR),

and fat retention (FR)

of *Channa striata* reared in the different mediums during the study

Parameters Treatments Ameliorant Lime (Dolomite) Control SR (%) 70.55 ± 7.21 a 62.58 ± 3.60 a 54.45 ± 3.13 b SGR (% day⁻¹) 6.42 ± 0.33 a 5.54 ± 0.21 b 4.77 ± 0.39 c Albumin content (g per 100 mL) 3.12 ± 0.06 a 2.88 ± 0.10 b 2.69 ± 0.08 c Feed efficiency (%) 89.07 ± 6.33 a 79.66 ± 6.98 a 67.61 ± 6.36 b PR (%) 53.41 ± 4.70 a 47.98 ± 4.48 a 40.79 ± 3.14 b FR (%) 89.41 ± 13.17 a 88.04 ± 14.67 a 76.79 ± 8.03 a

Different superscript letters in the same row indicate significant

different results ($p > 0.05$). *C. striata* is classified as a fish that is able to live in marginal medium, but it is sensitive to extreme environmental changes as well as majority of organisms. In relation to environmental conditions, the survival of the fish will be high, if the living medium is in a good condition, but when the living medium is not in a good condition, it will lead to the increasing of the fish mortality. The survivals of *C. striata* juveniles in the current results using ameliorant and lime were higher than the previous studies conducted in the laboratory using acidic sulfate water medium that resulted SR at a value of 54.44% (Purnamawati et al 2017a), but those were lower than the nursery phase in soil ponds with an SR of 57.33–76.67% (Rahman et al 2013) and the rearing in acidic sulfate water medium without aeration (SR 92%) (Purnamawati et al 2017b). The administration of ameliorant is able to produce water quality conditions that are appropriate to the habitat of *C. striata*. The improvement of water quality in acidic sulfate water through the addition of an ameliorant allows the fish to utilize feed more

AAFL Bioflux, 2018, Volume 11, Issue 6. 1923 <http://www.bioflux.com.ro/aafl>

optimal. The

pH range of 6.8–7.4 in the ameliorant and lime treatments is still good enough for the growth of *C. striata* juveniles, because it is still in tolerance range of pH for *C. striata* life (4.25–9.4) (Courtenay & Williams 2004), while the optimal pH for the intensive fish farming is 7.0–8.0 (Wedemeyer 1996). In this pH range, gills can maximally bind oxygen and ions required by the body, so that the physiological activities of the fish will be more optimal. The use of ameliorant is related to the increase in Ca^{2+} needed to bind sulfate (SO_4^{2-}) (a source of the water acidity) to form gypsum deposits (Fitzpatrick et al 1998). Humic acid, fulvic acid and humin acid contained in an ameliorant cause an increase in pH (H^+ decreases). The sulfate level in ameliorant treatment was 10.2–12.6 mg L⁻¹, this range was still within the tolerance range for the fish life. In all treatments, the sulfate levels in the experimental mediums were low and still within the tolerance range for fish life according to Boyd (1988). The improvement of water quality in ponds located in acid sulfate land through the administration of an ameliorant produced from banana stems in *C. striata* farming provided the best results compared to the control and a treatment using lime, because the water's toxicity potential became reduced. This was shown by a lower ammonia (NH_3) level (0.01–0.05 mg L⁻¹) in ameliorant treatment compared to other treatments. This level can still be tolerated by *C. striata*, because

at ammonia level of 1.57 mg L^{-1} , *C. striata* juvenile can still grow (Qin et al 1997).

Dissolved oxygen (DO) level range during the study was $4.12\text{-}5.80 \text{ mg L}^{-1}$. This range still meets the range recommended for the life and the growth of *C. striata* (Boyd 1988). *C. striata* is categorized in air breathing fish that has diverticula as the additional respiratory organ, so the fish can live in poor-oxygen-medium and can tolerate hypoxia by using oxygen derived from the air (Huang et al 2011; Song et al 2013). Alkalinity in water is caused by bicarbonate (HCO_3^-), carbonate (CO_3^{2-}), and hydroxide (OH^-) ions. At the ameliorant treatment, the alkalinity level was $85.1\text{-}115.8 \text{ mg L}^{-1} \text{ CaCO}_3$. This level was slightly higher than those of the lime treatment and control, and it was not far from the optimal range ($100\text{-}150 \text{ mg L}^{-1} \text{ CaCO}_3$) (Wedemeyer 1996). In the optimal range, the ability of alkalinity to sustain the pH of the water becomes better, but it does not negatively affect the fish growth (Wedemeyer 1996; Boyd 1988). The results of statistical analysis showed that the administration of an ameliorant and lime significantly affected the survival (SR) of *C. striata*. The administration of an ameliorant at a dose of 300 kg per ha pond and lime at a dose of 180 kg per ha pond produced a higher SR compared to control ($p > 0.05$). This was due to the better water quality in both treatment mediums compared to control. This caused the stress level of the fish decreasing and the fish became healthier. As a consequence, SR in ameliorant and lime treatments was also higher. Growth is a complex biological process. Growth can occur when there is excess energy and materials derived from the feed. The *C. striata* juveniles which were reared in the ameliorant treated pond produced the highest specific growth rate (SGR) (6.42%), followed by the lime and the control groups with SGR values of 5.54% and 4.77%, respectively ($p > 0.05$) (Table 2). The difference in growth rates showed that the *C. striata* which were reared in the ameliorant treated pond were better in utilizing the energy sourced by the feed. The high SGR in *C. striata* juveniles reared in the ameliorant treated pond was supported by a finding showing the highest albumin content in the fish blood (3.12 g per 100 mL) found in ameliorant treatment compared to other treatments ($p > 0.05$). A better water quality in the ameliorant treated pond will improve the ability of the fish to absorb nutrient from the feed. It is reflected in the high albumin content in the fish reared in the ameliorant treated pond. By maintaining albumin in the blood plasma, the fish can also maintain blood volume. Albumin is one of blood plasma proteins synthesized in the liver (Infusino & Panteghini 2013). Albumin plays a role in transporting small molecules through plasma and extracellular fluid (Suprayitno 2014), maintaining plasma balance in blood vessels with fluid in the interstitial cavity and the formation of cells and tissues (Nugroho 2012).

AACL Bioflux, 2018, Volume 11, Issue 6. 1924 <http://www.bioflux.com.ro/aac1>

C. striata juveniles which were reared in the medium treated with an ameliorant had higher feed intake and feed efficiency than those of control, but those were not significantly different from the lime treatment. This showed that the feed given could be properly utilized by the fish, so that nutrients contained in the feed could be retained in the fish body efficiently. The feed efficiency is determined by several factors such as genetic, nutritional content, farming management, and environment (Robinson & Li 2015; Hasan & Soto 2017). In normal conditions, physiological processes in the fish's body will run well, so that the feed

can be used entirely by the fish and can be converted into the fish flesh (Nelson & Chabot 2011). Daily feed intake

is one of factors that influence the potential of the fish to grow optimally and feed consumption rate is closely related to the capacity and the emptying of the

fish gut (Abowei & Ekubo 2011). The high SGR in the ameliorant treatment was not only reflected in the albumin content and feed efficiency, but also reflected in protein retention (PR). The ameliorant and lime treatments resulted in higher PR (53.41% and 47.98%) than that of control (Table 2). Protein retention is affected by the type of the feed, feed quality and environmental conditions (Samantaray & Mohanty 1997; Purnamawati et al 2017a). The better is the feed quality and the environment, higher is the protein retention. The administration of an ameliorant and lime into the pond has a major role in improving water quality, so that the utilization of the feed energy becomes more optimal. This causes the protein derived from the feed will be retained more in the body

and decomposed only in a few amount or it will be utilized as energy to maintain homeostasis.

Jobling et al (1991) state that the high energy for body activities will reduce the energy budget for the growth. According to Nelson & Chabot (2011), energy in the feed is physiologically used for activities and metabolism,

while the residue will be deposited as body tissue through the growth process and the synthesis of reproductive tissue. The difference in water quality conditions as a result of different treatments did not significantly influence fat retention ($p < 0.05$). This described that the utilization of fat by the fish in three experimental groups was not much different. This is thought to occur because the energy derived from the feed, especially derived from protein, is widely released by the body for the adaptation to poor environmental conditions.

Conclusions. An ameliorant from banana stems can be used as a substitute for lime in improving water quality of a pond located in acid sulfate land. The use of an ameliorant and dolomite resulted in higher survivals of the *C. striata* than that of without using an ameliorant and dolomite, but the best growth was obtained in the ameliorant group.

References

Abowei J. F. N., Ekubo E. T., 2011 Some principles and requirements in fish nutrition. *British Journal of Pharmacology and Toxicology* 2:163-178.

Alwi M., 2014 [The prospect of tidal swamp land for rice crops]. *The Proceeding of the National Seminar "Location-Specific Technological Innovation"*, 2014 August 6-7; Banjarbaru, Indonesia, pp. 45-59. [In Indonesian]. Arsyad D. M., Saidi B. B., Enrizal, 2014 [Development of agricultural innovations in tidal swamp land for increasing food sovereignty]. *Pengembangan Inovasi Pertanian* 7:169-178. [In Indonesian].

Boyd C. E., 1988 *Water quality in warmwater fish ponds*. Alabama Experiment Station, Auburn, 359

p. Cavalcante D. H., da Silva S. R., Pinheiro P. D., Akao M. M. F., e Sá M. V. C., 2012 Single or paired increase of total alkalinity and hardness of water for cultivation of Nile tilapia juveniles, *Oreochromis niloticus*. *Acta Scientiarum* 34:177-183.

Courtenay Jr.

W. R., Williams J. D., 2004 Snakeheads (Pisces, Channidae) - A biological synopsis and risk assesment. U.S. Geological Survey,

Denver, 143 p.

AAFL Bioflux, 2018, Volume 11, Issue 6. 1925 <http://www.bioflux.com.ro/aafl>

de

Queiroz J. F., Nicolella G., Wood C. W., Boyd C. E., 2004 Lime application methods, water and bottom soil acidity in fresh water fish ponds. *Scientia Agricola* 61:469- 475. Fitzpatrick R. W., Merry R. H., Williams J., White I., Bowman G., Taylor G., 1998 Acid sulfate soil assessment: coastal, inland and minespoil conditions. National Land and Water Resources Audit Methods Paper. National Land and Water Resources Department, Canberra, 18 p. Gam L. H., Leow C. Y., Baie S., 2006

Proteomic analysis of snakehead fish (*Channa striata*) muscle tissue. *Malaysian Journal of Biochemistry and Molecular Biology* 14:25-32.

Hasan M. R., Soto D., 2017 Improving feed conversion ratio and its impact on reducing greenhouse gas emissions in aquaculture. FAO Non-Serial Publication, Food and Agriculture Organization of the United Nations, Rome, 33 p.

Hidayatullah S., Muslim, Taqwa F. H., 2015 [Rearing of snakehead larvae (*Channa striata*) in plastic lined pond with different stocking density]. *Jurnal Perikanan dan Kelautan* 20:61-70. [In Indonesian].

Huang C. Y., Lin C. P., Lin H. C., 2011 Morphological and biochemical variations in the gills of 12 aquatic air-breathing anabantoid fish. *Physiological and Biochemical Zoology* 84:125-134.

Infusino I., Panteghini M., 2013 Serum albumin: accuracy and clinical use. *Clinica Chimica Acta* 419:15-18.

Jobling M., Knudsen R., Pedersen P. S., dos Santos J., 1991 Effects of dietary composition and energy content on the nutritional energetics of cod, *Gadus morhua*. *Aquaculture* 92:243-257.

Kang'ombe J., Brown J. A., 2008 Effect of salinity on growth, feed utilization, and survival of *Tilapia rendalli* under laboratory conditions. *Journal of Applied Aquaculture* 20:256-271.

Marimuthu K., Jesu Arokiaraj A., Haniffa M. A., 2009 Effect of diet quality on seed production of the spotted snakehead *Channa punctatus* (Bloch). *American-Eurasian Journal of Sustainable Agriculture* 3:344-347.

Mohd S. M. A., Abdul M. M. J., 2012

Therapeutic potential of the haruan (*Channa striatus*): from food to medicinal uses.

Malaysian Journal of Nutrition 18:125–136.

Mollah M. F. A., 1985 Effects of stocking density and water depth on the growth and survival of freshwater catfish (*Clarias macrocephalus* Gunther) larvae. Indian Journal of Fisheries 35:1–17.

Mollah M. F. A., Mamun M. S. A., Sarowar M. N., Roy A., 2009 Effects of stocking density on the growth and breeding performance of broodfish and larval growth and survival of shol, *Channa striatus* (Bloch). Journal of the Bangladesh Agricultural University 7:427-432.

Nakkrasae L., Wisetdee K., Charoenphandhu N., 2015

Osmoregulatory adaptations of freshwater air-breathing snakehead fish (*Channa striata*) after exposure to brackish water.

Journal of Comparative Physiology B 185:527–537.

Nelson J. A., Chabot D., 2011 General energy metabolism. In: Encyclopedia of fish physiology: from genome to environment, volume 3. Farrell A. P. (ed), pp. 1566- 1572, Academic Press, San Diego.

Nugroho M., 2012 [The isolation of albumin and characteristics of molecular weight from extraction result by steaming of fish gabus (*Ophiocephalus striatus*)]. SAINTEK PERIKANAN: Indonesian Journal of Fisheries Science and Technology 9:40–48. [In Indonesian].

Purnamawati, 2017 [The growth performance of snakehead *Channa striata* Bloch. on tidal land through water quality engineering]. Bogor Agricultural University, Bogor, 78 p. [In Indonesian].

Purnamawati, Djokosetiyanto D., Nirmala K., Harris E., Affandi R., 2017a

Survival and growth of striped snakehead fish (*Channa striata* Bloch.) juvenile reared in acid sulfate water and rainwater medium. AACL Bioflux 10:265–273.

Purnamawati, Djokosetiyanto D., Nirmala K., Surawidjaja E. H., Affandi R., 2017b

Survival and growth response of snakehead fish *Channa striata* Bloch juvenile

in

AACL Bioflux, 2018, Volume 11, Issue 6. 1926 <http://www.bioflux.com.ro/aac>

aerated and unaerated acid sulfate water.

Jurnal Akuakultur Indonesia 16:60–67.

Qin J.,

Fast A. W., Kai A. T., 1997 Tolerance of snakehead *Channa striatus* to ammonia at different pH. *Journal of the World Aquaculture Society* 28:87-90.

Rahman M. A., Arshad A.,

Amin S. M. N., Shamsudin M. N., 2013 Growth and survival of fingerlings of a threatened snakehead, *Channa striatus* (Bloch) in earthen nursery ponds. *Asian Journal of Animal and Veterinary Advances* 8:216–226. Robinson E. H., Li M. H., 2015 Feed conversion ratio for pond-raised catfish. Information Sheet No. 1364. The Office of Agricultural Communications, The

Division of Agriculture, Forestry, and Veterinary Medicine, Mississippi State University, Strakville, 4 p.

Samantaray K., Mohanty S. S., 1997

Interactions of dietary levels of protein and energy on fingerling snakehead, *Channa striata*. *Aquaculture* 156:241–249.

Song L. M., Munian K., Rashid Z. A., Bhassu S., 2013 Characterization

of Asian snakehead murrel *Channa striata* (Channidae) in Malaysia: an insight into molecular data and morphological approach.

The

Scientific World Journal 2013, Article ID 917506, <http://dx.doi.org/10.1155/2013/917506>.

Suprayitno E., 2014 Profile albumin fish cork (*Ophicephalus striatus*) of different ecosystems. *International Journal of Current Research and Academic Review* 2:201– 208.

Takeuchi T., 1988 Laboratory work-chemical evaluation of dietary nutrient. In: *Fish nutrition and mariculture*. Watanabe T. (ed), pp. 179-233, Kanagawa International Fisheries Training Centre, Japan International Cooperation Agency (JICA), Tokyo. Vivekanandan E., 1977 Surfacing activity and food utilization in the obligatory air-breathing fish *Ophiocephalus striatus* as a function of body weight. *Hydrobiologia* 55:99-112.

Weatherley A. H., Gill H. S., 1989 *The biology of fish growth*. Academic Press, London, 443 p. Wedemeyer G. A., 1996 *Physiology of fish in intensive culture system*. Springer US, New York, 232 p.

Wedemeyer G. A., Yasutake

W. T., 1977

Clinical methods for the assessment of

the effects of environmental stress on fish health.

U.S. Fish

and Wildlife Service,

Washington D.C., 18

p. ***

APHA (

American Public Health Association), 1989

Standard methods for the examination of water and wastewater 17 th

edition.

American Public Health Association,

Washington D.C., 1268

p. *** BPS Kalbar (Badan Pusat Statistik Provinsi Kalimantan Barat), 2012 [West Kalimantan in number]. Badan Pusat Statistik Provinsi Kalimantan Barat, Pontianak. [In Indonesian].

Received: 01 November 2018. Accepted: 02 December 2018. Published online: 10 December 2018. Authors: Purnamawati,

Pontianak State Polytechnic, Department of Marine Science and Fisheries, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: pur_polnep@yahoo.com Nurmalia, Pontianak State Polytechnic, Department of Marine Science and Fisheries, Study Program of Aquaculture, Indonesia, West

Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: ibu.nurmalia@gmail.com Mochammad Idham Shilman,

Pontianak State Polytechnic, Department of Marine Science and Fisheries, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail:

idham.shilman@yahoo.co.id Eko Dewantoro, Muhammadiyah University of Pontianak, Faculty of Fisheries and

Marine Science, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: ekodewe_ump@yahoo.com

Diah Ayu Satyari Utami, Marine and Fisheries Polytechnic of Jembrana, Study Program of Aquaculture, Indonesia, Bali, Jembrana 82218, Desa Pengambangan,

e-mail: dplongitm@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited. How to cite this article: Purnamawati, Nurmala, Shilman M. I., Dewantoro E., Utami D. A. S., 2018 Effects of ameliorant on the growth of snakehead fish (*Channa striata*) juvenile reared in acid sulfate water medium located in tidal land. *AAFL Bioflux* 11(6):1919-1926.

Hit and source - focused comparison, Side by Side:

Left side: As student entered the text in the submitted document.

Right side: As the text appears in the source.

Instances from: <http://www.bioflux.com.ro/docs/2018.1919-1926.pdf>

1 100%

AAFL Bioflux, 2018, Volume 11, Issue 6. 1919 <http://www.bioflux.com.ro/aafl>

Effects of ameliorant on the growth of snakehead fish (*Channa striata*) juvenile reared in acid sulfate water

medium located in tidal land 1 Purnamawati, 1 Nurmala, 1 Mochammad I. Shilman, 2 Eko Dewantoro, 3 Diah A. S. Utami 1

Study Program of Aquaculture, Department of Marine Science and Fisheries, Pontianak State Polytechnic, Pontianak, West Kalimantan, Indonesia; 2 Study Program of Aquaculture, Faculty of Fisheries and Marine Science,

Muhammadiyah University of Pontianak, Pontianak, West Kalimantan, Indonesia; 3 Study Program of Aquaculture, Marine and Fisheries Polytechnic of Jembrana, Jembrana, Bali,

1: <http://www.bioflux.com.ro/docs/2018.1919-1926.pdf> 100%

AAFL Bioflux, 2018, Volume 11, Issue 6. 1919 <http://www.bioflux.com.ro/aafl> Effects of ameliorant on the growth of snakehead fish (*Channa striata*) juvenile reared in acid sulfate water medium located in tidal land 1 Purnamawati, 1 Nurmala, 1 Mochammad I. Shilman, 2 Eko Dewantoro, 3 Diah A. S. Utami 1 Study Program of Aquaculture, Department of Marine Science and Fisheries, Pontianak State Polytechnic, Pontianak, West Kalimantan, Indonesia; 2 Study Program of Aquaculture, Faculty of Fisheries and Marine Science, Muhammadiyah University of Pontianak, Pontianak, West Kalimantan, Indonesia; 3 Study Program of Aquaculture, Marine and Fisheries Polytechnic of Jembrana, Jembrana, Bali, Indonesia. Corresponding author: Purnamawati, pur_polnep@yahoo.com Abstract. The aim of this study was to evaluate the effects of the use of an ameliorant produced from banana stems to increase water pH and improve the growth of the snakehead fish (*Channa striata*). The experiment applied a completely randomized design. The

Indonesia. Corresponding author: Purnamawati, pur_polnep@yahoo.com Abstract. The aim of this study was to evaluate the effects of the use of

an ameliorant produced from banana stems to increase water pH and improve the growth of the snakehead fish (*Channa striata*). The experiment applied

a completely randomized design. The treatments applied in this study consisted of three treatments

including ameliorant (300 kg ha⁻¹), lime (CaCO₃) (180 kg ha⁻¹), and control (without ameliorant and lime). Each treatment applied four replicates.

C. striata juveniles with an average initial length of 2.5±0.2 cm and an average initial weight of 0.27±0.03 g were reared in 12 units of hapa net cages measuring 2 x 3 x 1.5 m with a stocking density of 2 fish per L, for 60 days. The fish were fed commercial feed with a

protein content of 40%,

twice a day (morning and afternoon) until to apparent satiation. The

use of an ameliorant in the experimental medium significantly gave better

treatments applied in this study consisted of three treatments including ameliorant (300 kg ha⁻¹), lime (CaCO₃) (180 kg ha⁻¹), and control (without ameliorant and lime). Each treatment applied four replicates. *C. striata* juveniles with an average initial length of 2.5±0.2 cm and an average initial weight of 0.27±0.03 g were reared in 12 units of hapa net cages measuring 2 x 3 x 1.5 m with a stocking density of 2 fish per L, for 60 days. The fish were fed commercial feed with a protein content of 40%, twice a day (morning and afternoon) until to apparent satiation. The use of an ameliorant in the experimental medium significantly gave better results shown by higher survival (70.55%), specific growth rate (6.42% day⁻¹), feed efficiency (89.07%), protein retention (53.41%), fat retention (89.41%), and albumin content (3.12 g per 100 mL) than those of control. Key Words: air-breathing fish, banana stems, dolomite, tidal area. Introduction. Indonesia has about 20.11 million ha of tidal swamp land consisting of 2.07 million ha of tidal potential land, 6.71 million ha of acid sulfate land, 10.89 million ha of peatland, and 0.44 million ha of saline land. The tidal land is spread in several regions, including Sumatera, Kalimantan, Papua, and Sulawesi (Arsyad et al 2014). Tidal swamp land area of West Kalimantan is around 1.12 million ha, which is spread in several districts (BPS Kalbar 2012) and 20,532 ha of it is acid sulfate land (Alwi 2014). Most of these lands have been used for fish farming, but in general, the productivity is still relatively low. The low productivity of fisheries sectors in this area is due to various problems, especially poor water quality. In a low water quality condition, there are only certain fish species that can be cultivated, one of those species is

results shown by higher survival (70.55%), specific growth rate (6.42% day⁻¹), feed efficiency (89.07%), protein retention (53.41%), fat retention (89.41%),

and albumin content (3.12 g per 100 mL) than those of control.

Key Words: air-breathing fish, banana stems, dolomite, tidal area. Introduction. Indonesia has about 20.11 million ha of tidal swamp land consisting of 2.07 million ha of tidal potential land, 6.71 million ha of acid sulfate land, 10.89 million ha of peatland, and 0.44 million ha of saline land. The tidal land is spread in several regions, including Sumatera, Kalimantan, Papua, and Sulawesi (Arsyad et al 2014). Tidal swamp land area of West Kalimantan is around 1.12 million ha, which is spread in several districts (BPS Kalbar 2012) and 20,532 ha of it is acid sulfate land (Alwi 2014).

Most of these lands have been used for fish farming, but in general, the productivity is still relatively low. The low productivity

of fisheries sectors in this area is due to various problems, especially poor water quality. In a low water quality condition, there are only certain fish species that can be cultivated, one of those species is the snakehead fish (*Channa striata*). *C. striata*, is a freshwater fish species, classified in Channidae family and Perciformes order (Nakkrasae et al 2015). This fish is a potential and important species to be developed as an aquaculture commodity (Mollah 1985; Marimuthu et al 2009; Mollah et al 2009; Rahman et al 2013), and has high economic value with

the snakehead fish (*Channa striata*). *C. striata*, is a freshwater fish species, classified in Channidae family and Perciformes order (Nakkrasae et al 2015). This fish is a potential and important species to be developed as an aquaculture commodity (Mollah 1985; Marimuthu et al 2009; Mollah et al 2009; Rahman et al 2013), and has high economic value with prices ranging from 3.78 to 4.47 USD kg⁻¹. Moreover, this fish flesh is used as a treatment of post- surgical therapy and can increase body endurance (Gam et al 2006; Marimuthu et al 2009; Mohd & Abdul 2012). *C. striata* usually lives in rivers, rice fields and ponds. *C. striata* relatively prefers calm water (without agitation). Another advantage possessed by *C. striata* is its ability to

AAFL Bioflux, 2018, Volume 11, Issue 6. 1920 <http://www.bioflux.com.ro/aafl> live in low oxygen environment. But to grow optimally, this species needs to live in a medium with optimal water quality ranges. One of limiting factors that influences the *C. striata* cultivation in acid sulfate land is pH. Tidal land has a pH range of 2.53–3.39 (Purnamawati et al 2017a), sulfate concentration at 6.91–8.7 mg L⁻¹, dissolved oxygen less than 5 mg L⁻¹ (Purnamawati et al 2017b). This water quality condition can be improved by increasing the pH of the culture medium. To increase the water pH in fish ponds can be done by adding a certain amount of lime into fish ponds (de Queiroz et al 2004; Cavalcante et al 2012). However, the required lime is not always available, especially in some locations that are far from the centers of agricultural production facilities. To overcome this issue, some local based plant materials can be

prices ranging from 3.78 to 4.47 USD kg⁻¹. Moreover, this fish flesh is used as a treatment of post- surgical therapy and can increase body endurance (Gam et al 2006; Marimuthu et al 2009; Mohd & Abdul 2012). *C. striata* usually lives in rivers, rice fields and ponds. *C. striata* relatively prefers calm water (without agitation). Another advantage possessed by *C. striata* is its ability to

AACL Bioflux, 2018, Volume 11, Issue 6. 1920 <http://www.bioflux.com.ro/aacl>

live in low oxygen environment. But to grow optimally, this species needs to live in a medium with optimal water quality ranges. One of limiting factors that influences the *C. striata* cultivation in acid sulfate land is pH. Tidal land has a pH range of 2.53–3.39 (Purnamawati et al 2017a), sulfate concentration at 6.91–8.7 mg L⁻¹, dissolved oxygen less than 5 mg L⁻¹ (Purnamawati et al 2017b). This water quality condition can be improved by increasing the pH of the culture medium. To increase the water pH in fish ponds can be done by adding a certain amount of lime into fish ponds (de Queiroz et al 2004; Cavalcante et al 2012). However, the required lime is not always available, especially in some locations that are far from the centers of agricultural production facilities. To overcome this issue, some local based plant materials can be used as ameliorants to substitute the utilization of lime. Banana stem is a waste that can be used as an ameliorant. This is caused by banana stem has some advantages, including low cost, abundant availability, easy to be processed, and it is not harmful

used as ameliorants to substitute the utilization of lime. Banana stem is a waste that can be used as an ameliorant. This is caused by banana stem has some advantages, including low cost, abundant availability, easy to be processed, and it is not harmful for the cultivated organisms. This material is expected to be a solution in aquaculture practices conducted in acid sulfate medium, especially in tidal land. This study aimed to evaluate the effectiveness of the use of banana stems as an ameliorant in acid sulfate medium in order to promote the growth of *C. striata* juveniles. Material and Method. The present study was conducted on July-September 2018. The rearing of *C. striata* was conducted in a tidal pond located in Kubu Raya District, West Kalimantan, Indonesia. The measurement of plasma glucose levels, albumin levels, the proximate analysis of the fish feed and the experimental fish were conducted at Environmental and Food Technology Laboratory, Tanjungpura University, Pontianak, Indonesia. Experimental design. This study was conducted through a completely randomized design (CRD) in a tidal pond using hapa net cages. The treatments applied in this study consisted of three treatments including ameliorant using banana stems (300 kg ha⁻¹), lime (CaCO₃) (180 kg ha⁻¹), and control (without ameliorant and lime). Each treatment applied four replicates. Experimental fish. The experimental fish used in this study were *C. striata* juveniles. The experimental fish had an average initial length of 2.5±0.2 cm and an average initial weight of 0.27±0.03 g. Experimental mediums. The experimental mediums used in this study were 12 units of hapa net cage with a size of 2 x 3 x 1.5 m. The hapa net cages were set in a pond, the

for the cultivated organisms. This material is expected to be a solution in aquaculture practices conducted in acid sulfate medium, especially in tidal land. This study aimed to evaluate the effectiveness of the use of banana stems as an ameliorant in acid sulfate medium in order to promote the growth of *C. striata* juveniles. Material and Method. The present

study was conducted on July-September 2018. The rearing of *C. striata* was conducted

in a tidal pond located in Kubu Raya District, West Kalimantan, Indonesia. The measurement of plasma glucose levels, albumin

levels, the proximate analysis of the fish feed and the experimental fish were

conducted

at Environmental and Food Technology Laboratory, Tanjungpura University, Pontianak, Indonesia. Experimental design. This study was conducted through a completely randomized design (CRD)

in a tidal pond using hapa net cages. The treatments applied in this study consisted of three treatments including ameliorant using banana stems (300 kg ha⁻¹), lime (CaCO₃) (180 kg ha⁻¹), and control (without ameliorant and lime).

Each treatment applied four replicates. Experimental fish. The experimental fish used in this study were *C. striata* juveniles.

top sides were covered with nets to avoid fish jumping out of the rearing cages. Experimental procedures. The production of ameliorant from banana stems used a bio-activator (EM4: effective microorganism 4) to accelerate the composting process. The composting process occurred aerobically. After the raw material became a compost, it was then dried under sunlight. The chemical composition of a compost from banana stems consisted of Ca (1.54%), Mg (0.86%), C-organic (43.39%), N (2.48%), C/N (17.50), pH (8.8), KTK + (41.07 cmol⁺ kg⁻¹), humic acid (2.50 ppm), fulvic acid (0.80 ppm), and moisture (68.78%) (Purnamawati 2017). Prior to the experiment, the experimental fish were acclimatized to experimental conditions. This step was carried out for 7 days in three net cages with a size of 4 x 3 x 1.5 m. *C. striata* were then reared in a tidal pond using hapa net cages with a stocking density of 2 fish per L (Vivekanandan 1977; Hidayatullah et al 2015). This study was conducted for 60 days. The experimental fish were fed commercial feed with a protein content of ±40%, twice a day to apparent satiation. Experimental parameters. The measurements of water quality parameters were performed following the procedures described by APHA (1989). The water quality parameters measured were temperature, pH, dissolved oxygen, sulfate (SO₄²⁻),

AAAL Bioflux, 2018, Volume 11, Issue 6. 1921 <http://www.bioflux.com.ro/aaal> alkalinity, and ammonia. The measurements of temperature, pH and dissolved oxygen level were carried out every 15 days, while the measurements of sulfate, alkalinity and ammonia were performed at the

The experimental fish had

an average initial length of 2.5 ± 0.2 cm and an average initial weight of 0.27 ± 0.03 g.

Experimental mediums. The experimental mediums used in this study were 12 units of hapa net cage with a size of 2 x 3 x 1.5 m. The hapa net cages were set

in a pond, the top sides were covered with nets to avoid fish jumping out of the rearing cages. Experimental procedures.

The production of ameliorant from banana stems used a bio-activator (EM4: effective microorganism 4) to accelerate the composting process. The composting process occurred aerobically. After the raw material became a compost, it was then dried under sunlight. The chemical composition of a compost from banana stems consisted of Ca (1.54%), Mg (0.86%), C-organic (43.39%), N (2.48%), C/N (17.50), pH (8.8), KTK + ($41.07 \text{ cmol} + \text{kg}^{-1}$), humic acid (2.50 ppm), fulvic acid (0.80 ppm), and moisture (68.78%) (Purnamawati 2017). Prior to the experiment, the experimental fish were acclimatized to experimental conditions. This step was carried out for 7 days in three net cages with a size of 4 x 3 x 1.5 m. *C. striata* were then reared in a tidal pond using hapa net cages with a stocking density of 2 fish per L (Vivekanandan 1977; Hidayatullah et al 2015). This study was conducted for 60 days. The experimental fish were fed commercial feed with a protein content of $\pm 40\%$, twice a day to apparent satiation.

beginning, middle and the end of the experiment. The measurement of albumin levels followed the method described by Infusino & Panteghini (2013), while the proximate analysis of the fish feed and the experimental fish were carried out according to the procedure by Takeuchi (1988). The measurement of plasma glucose levels was performed using a liquicolor glucose commercial kit GOD-PAP with calorimetric method and the results of the measurement were read with a spectrophotometer at a wavelength of 500 nm following the procedure constructed by Wedemeyer & Yasutake (1977). The daily mortality was recorded to obtain the survival data. Survival was calculated using a formula according to Kang'ombe & Brown (2008): $SR = (N_t \times N_0^{-1}) \times 100$ Note: SR = Survival (%) N_t = The number of the fish at the end of the study (individual) N_0 = The number of the fish at the beginning of the study (individual) The fish weight samplings were done every 15 days to obtain the growth data (specific growth rate). Specific growth rate was calculated using a formula stated by Weatherley & Gill (1989). $SGR = [(\ln W_2 - \ln W_1) / (t_2 - t_1)] \times 100$ Note: SGR = Specific growth rate (% day⁻¹) W_1 = The average weight of the fish at the beginning of the study (g) W_2 = The average weight of the fish at time t_2 (g) $t_2 - t_1$ = Experimental duration Feed intake (F) was obtained by summing the daily amount of the feed consumed during the study. The feed efficiency was calculated using a formula constructed by Kang'ombe & Brown (2008). $e = [(W_t + D) - W_o] / F \times 100$ Note: e = Feed efficiency W_o = The weight of the fish at the beginning of the study (g) W_t = The weight of the fish at the end of the study (g) D = The weight of

Experimental parameters. The measurements of water quality parameters were performed following the procedures described by APHA (1989). The

water quality parameters measured were temperature, pH, dissolved oxygen, sulfate (SO_4^{2-}),

alkalinity, and ammonia. The

AAFL Bioflux, 2018, Volume 11, Issue 6. 1921 <http://www.bioflux.com.ro/aac>

ammonia. The

measurements of temperature, pH and dissolved oxygen level were carried out every 15 days, while the measurements of sulfate, alkalinity and ammonia were performed at the beginning,

middle and the end of the experiment.

The measurement of albumin levels followed the method described by Infusino & Panteghini (2013), while the proximate analysis of the fish feed and the experimental fish were carried out according to the procedure by Takeuchi (1988). The measurement of plasma glucose levels was performed using a liquicolor glucose commercial kit GOD-PAP with calorimetric method and the results of the measurement were read with a spectrophotometer at a wavelength of 500 nm following the procedure

the dead fish during the study (g) F = The weight of total feed given during the study (g dry weight) Statistical analysis. The water quality parameters were analyzed through descriptive statistic. Survival, specific growth rate, albumin content, feed efficiency, protein retention, and fat retention were analyzed through ANOVA with a help of SPSS version 22 to see the differences among treatments. Results and Discussion. The water quality ranges in the experimental mediums are shown in Table 1. Water quality ranges varied among one treatment and other treatments. Water temperature ranges in the experimental mediums were not relatively different among treatments and those were an optimal range for the fish growth. The pH, total alkalinity, and sulfate showed differences between ameliorant added and lime added treatments to control (without the addition of ameliorant and lime). This could be seen from the high pH and total alkalinity, as well as the low sulfate levels dissolved in water. There was no difference in dissolved oxygen (DO) and ammonia levels among ameliorant, lime and control treatments.

AAFL Bioflux, 2018, Volume 11, Issue 6. 1922 <http://www.bioflux.com.ro/aac>

Table 1 The values of physical-chemical parameters of the water in each treatment during the study

Parameters	Treatments	Tolerance range *	optimum range **
pH	Ameliorant	6.8–7.2	6.9–7.4
	Lime (Dolomite)	5.3–5.8	4.25–9.4*
	Control	10.2–12.6	10.7–29.6
Sulfate (SO_4^{2-}) (mg L ⁻¹)	Ameliorant	20.1–48.2	5–150*
	Lime (Dolomite)	10.2–12.6	10.7–29.6
	Control	10.2–12.6	10.7–29.6
Alkalinity (mg L ⁻¹)	Ameliorant	85.1–115.8	71.0–110.4
	Lime (Dolomite)	22.1–26.6	100–150*
	Control	85.1–115.8	71.0–110.4
Ammonia (NH ₃) (mg L ⁻¹)	Ameliorant	0.01–0.05	
	Lime (Dolomite)	0.01–0.05	
	Control	0.01–0.05	

21

constructed by Wedemeyer & Yasutake (1977). The daily mortality was recorded to obtain the survival data.

Survival was calculated using a formula according to Kang'ombe & Brown (2008): $SR = (N_t \times N_0^{-1}) \times 100$ Note: SR = Survival (%) N_t = The number of the fish at the end of the study (individual) N_0 =

The number of the fish at the beginning of the study (

individual) The fish weight samplings were done every 15 days to obtain the growth data (specific growth rate).

Specific growth rate was calculated using a formula stated by Weatherley & Gill (1989). $SGR = [(\ln W_2 - \ln W_1) / (t_2 - t_1)] \times 100$

Note:

SGR = Specific growth rate (% day⁻¹) W_1 = The average weight of the fish at the beginning of the study (g) W_2 = The average weight of the fish at time t_2 (g) $t_2 - t_1$ = Experimental duration
Feed

intake (F) was obtained by summing the daily amount of the feed consumed during the study. The feed efficiency

was calculated using a formula constructed by Kang'ombe & Brown (2008). $e = [(W_t + D) - W_0] / F \times 100$

0.01–0.07 0.03–0.05 0.54–1.57* 2) DO (mg L⁻¹) 4.12–5.80
4.15–5.47 4.79–5.12 < 5 3) Temperature (°C) 29.32–30.12
29.42–30.29 28.32–30.02 26–32** 3) 1) Wedemeyer (1996), 2 Qin
et al (1997), 3) Courtenay, Jr. & Williams (2004), *) tolerance
range, **) optimum range. The growth performance (survival,
SGR and feed efficiency) and physiological characteristics
(protein retention, energy retention, and albumin content) of *C.*
striata in 60-day-experiment can be seen in Table 2. The addition
of an ameliorant into ponds located in an acid sulfate land could
produce the highest specific growth rate (SGR) and albumin level
of *C. striata* (6.42% day⁻¹ and 3.12 g per 100 mL, respectively),
followed by lime added and control treatments ($p > 0.05$). The
survival (SR) of *C. striata* juveniles ranged from 54.45 to 70.55%.
The highest SR was obtained in ameliorant treatment, but it was
not significantly different from the lime added treatment. The
same results were also found in feed efficiency and protein
retention (PR), while fat retention (FR) does not show a
significant difference between among treatments tested in this
study. Table 2 Survival (SR), specific growth rate (SGR), albumin
content, feed efficiency, protein retention (PR), and fat retention
(FR) of *Channa striata* reared in the different mediums during
the study Parameters Treatments Ameliorant Lime (Dolomite)
Control SR (%) 70.55±7.21 a 62.58±3.60 a 54.45±3.13 b SGR (%
day⁻¹) 6.42±0.33 a 5.54±0.21 b 4.77±0.39 c Albumin content (g
per 100 mL) 3.12±0.06 a 2.88±0.10 b 2.69±0.08 c Feed efficiency
(%) 89.07±6.33 a 79.66±6.98 a 67.61±6.36 b PR (%) 53.41±4.70 a
47.98±4.48 a 40.79±3.14 b FR (%) 89.41±13.17 a 88.04±14.67 a
76.79±8.03 a Different superscript letters in the same row

Note: e =

Feed efficiency W_o = The weight of the fish at the beginning of the study (g) W_t = The weight of the fish at the end of the study (g) D = The weight of the dead fish during the study (g) F = The weight of total feed given during the study (g dry weight)

Statistical analysis. The water quality parameters were analyzed through descriptive statistic. Survival, specific growth rate, albumin content, feed efficiency, protein retention,

and fat retention were analyzed through ANOVA with a help of SPSS version 22

to see the differences among treatments. Results and Discussion. The

water quality ranges in the experimental mediums are shown in Table 1. Water quality ranges varied among one treatment and other treatments. Water temperature ranges in the experimental mediums were not relatively different among treatments and those were an optimal range for the fish growth. The pH, total alkalinity, and sulfate showed differences between ameliorant added and lime added treatments to control (without the addition of ameliorant and lime). This could be seen from the high pH and total alkalinity, as well as the low sulfate levels dissolved in water. There was no difference in dissolved oxygen (DO) and ammonia levels among ameliorant, lime and control treatments.

indicate significant different results ($p > 0.05$). *C. striata* is classified as a fish that is able to live in marginal medium, but it is sensitive to extreme environmental changes as well as majority of organisms. In relation to environmental conditions, the survival of the fish will be high, if the living medium is in a good condition, but when the living medium is not in a good condition, it will lead to the increasing of the fish mortality. The survivals of *C. striata* juveniles in the current results using ameliorant and lime were higher than the previous studies conducted in the laboratory using acidic sulfate water medium that resulted SR at a value of 54.44% (Purnamawati et al 2017a), but those were lower than the nursery phase in soil ponds with an SR of 57.33–76.67% (Rahman et al 2013) and the rearing in acidic sulfate water medium without aeration (SR 92%) (Purnamawati et al 2017b). The administration of ameliorant is able to produce water quality conditions that are appropriate to the habitat of *C. striata*. The improvement of water quality in acidic sulfate water through the addition of an ameliorant allows the fish to utilize feed more

AAFL Bioflux, 2018, Volume 11, Issue 6. 1923 <http://www.bioflux.com.ro/aafl> optimal. The pH range of 6.8-7.4 in the ameliorant and lime treatments is still good enough for the growth of *C. striata* juveniles, because it is still in tolerance range of pH for *C. striata* life (4.25-9.4) (Courtenay & Williams 2004), while the optimal pH for the intensive fish farming is 7.0–8.0 (Wedemeyer 1996). In this pH range, gills can maximally bind oxygen and ions required by the body, so that the physiological

AACL

Bioflux, 2018, Volume 11, Issue 6. 1922 <http://www.bioflux.com.ro/aac>

Table 1

The values of physical-chemical parameters of the water in each treatment during the study Parameters Treatments Tolerance range * and optimum range ** Ameliorant Lime (Dolomite) Control pH 6.8–7.2 6.9–7.4 5.3–5.8 4.25–9.4* 3) Sulfate (SO₄²⁻) (mg L⁻¹) 10.2–12.6 10.7–29.6 20.1–48.2 5–150* 1) Alkalinity (mg L⁻¹) 85.1–115.8 71.0–110.4 22.1–26.6 100–150* 1) Ammonia (NH₃) (mg L⁻¹) 0.01–0.05 0.01–0.07 0.03–0.05 0.54–1.57* 2) DO (mg L⁻¹) 4.12–5.80 4.15–5.47 4.79–5.12 < 5 3) Temperature (°C) 29.32–30.12 29.42–30.29 28.32–30.02 26–32** 3) 1) Wedemeyer (1996), 2) Qin et al (1997), 3) Courtenay, Jr. & Williams (2004), * tolerance range, ** optimum range.

The growth performance (survival, SGR and feed efficiency) and physiological characteristics (protein retention, energy retention, and albumin content) of *C. striata* in 60-day-experiment can be seen in Table 2. The addition of an ameliorant into ponds located in an acid sulfate land could produce the highest specific growth rate (SGR) and albumin level of *C. striata* (6.42% day⁻¹ and 3.12 g per 100 mL, respectively), followed by lime added and control treatments ($p > 0.05$). The survival (SR) of *C. striata* juveniles ranged from 54.45 to 70.55%. The highest SR was obtained in ameliorant treatment, but it was not significantly different from

activities of the fish will be more optimal. The use of ameliorant is related to the increase in Ca²⁺ needed to bind sulfate (SO₄²⁻) (a source of the water acidity) to form gypsum deposits (Fitzpatrick et al 1998). Humic acid, fulvic acid and humin acid contained in an ameliorant cause an increase in pH (H⁺ decreases). The sulfate level in ameliorant treatment was 10.2–12.6 mg L⁻¹, this range was still within the tolerance range for the fish life. In all treatments, the sulfate levels in the experimental mediums were low and still within the tolerance range for fish life according to Boyd (1988). The improvement of water quality in ponds located in acid sulfate land through the administration of an ameliorant produced from banana stems in *C. striata* farming provided the best results compared to the control and a treatment using lime, because the water's toxicity potential became reduced. This was shown by a lower ammonia (NH₃) level (0.01–0.05 mg L⁻¹) in ameliorant treatment compared to other treatments. This level can still be tolerated by *C. striata*, because at ammonia level of 1.57 mg L⁻¹, *C. striata* juvenile can still grow (Qin et al 1997). Dissolved oxygen (DO) level range during the study was 4.12–5.80 mg L⁻¹. This range still meets the range recommended for the life and the growth of *C. striata* (Boyd 1988). *C. striata* is categorized in air breathing fish that has diverticula as the additional respiratory organ, so the fish can live in poor-oxygen-medium and can tolerate hypoxia by using oxygen derived from the air (Huang et al 2011; Song et al 2013). Alkalinity in water is caused by bicarbonate (HCO₃⁻), carbonate (CO₃²⁻), and hydroxide (OH⁻) ions. At the ameliorant treatment, the alkalinity level was 85.1–115.8 mg L⁻¹

the lime added treatment. The same results were also found in feed efficiency and protein retention (PR), while fat retention (FR) does not show a significant difference between among treatments tested in this study.

Table 2 Survival (SR), specific growth rate (SGR), albumin content, feed efficiency, protein retention (PR),

and fat retention (FR)

of *Channa striata* reared in the different mediums during the study

Parameters	Treatments	Ameliorant	Lime (Dolomite)	Control	SR (%)	
SR (%)	70.55±7.21	a	62.58±3.60	a	54.45±3.13	b
SGR (% day ⁻¹)	6.42±0.33	a	5.54±0.21	b	4.77±0.39	c
Albumin content (g per 100 mL)	3.12±0.06	a	2.88±0.10	b	2.69±0.08	c
Feed efficiency (%)	89.07±6.33	a	79.66±6.98	a	67.61±6.36	b
PR (%)	53.41±4.70	a	47.98±4.48	a	40.79±3.14	b
FR (%)	89.41±13.17	a	88.04±14.67	a	76.79±8.03	a

Different superscript letters in the same row indicate significant

different results ($p > 0.05$). *C. striata* is classified as a fish that is able to live in marginal medium, but it is sensitive to extreme environmental changes as well as majority of organisms. In relation to environmental conditions, the survival of the fish will be high, if the living medium is in a good condition, but when the living medium is not in a good condition, it will lead to the increasing of the fish mortality. The survivals of *C. striata* juveniles in the current results using ameliorant and lime were

CaCO_3 . This level was slightly higher than those of the lime treatment and control, and it was not far from the optimal range (100-150 mg L⁻¹ CaCO_3) (Wedemeyer 1996). In the optimal range, the ability of alkalinity to sustain the pH of the water becomes better, but it does not negatively affect the fish growth (Wedemeyer 1996; Boyd 1988). The results of statistical analysis showed that the administration of an ameliorant and lime significantly affected the survival (SR) of *C. striata*. The administration of an ameliorant at a dose of 300 kg per ha pond and lime at a dose of 180 kg per ha pond produced a higher SR compared to control ($p > 0.05$). This was due to the better water quality in both treatment mediums compared to control. This caused the stress level of the fish decreasing and the fish became healthier. As a consequence, SR in ameliorant and lime treatments was also higher. Growth is a complex biological process. Growth can occur when there is excess energy and materials derived from the feed. The *C. striata* juveniles which were reared in the ameliorant treated pond produced the highest specific growth rate (SGR) (6.42%), followed by the lime and the control groups with SGR values of 5.54% and 4.77%, respectively ($p > 0.05$) (Table 2). The difference in growth rates showed that the *C. striata* which were reared in the ameliorant treated pond were better in utilizing the energy sourced by the feed. The high SGR in *C. striata* juveniles reared in the ameliorant treated pond was supported by a finding showing the highest albumin content in the fish blood (3.12 g per 100 mL) found in ameliorant treatment compared to other treatments ($p > 0.05$). A better water quality in the ameliorant treated pond will improve

higher than the previous studies conducted in the laboratory using acidic sulfate water medium that resulted SR at a value of 54.44% (Purnamawati et al 2017a), but those were lower than the nursery phase in soil ponds with an SR of 57.33–76.67% (Rahman et al 2013) and the rearing in acidic sulfate water medium without aeration (SR 92%) (Purnamawati et al 2017b). The administration of ameliorant is able to produce water quality conditions that are appropriate to the habitat of *C. striata*. The improvement of water quality in acidic sulfate water through the addition of an ameliorant allows the fish to utilize feed more

AAFL Bioflux, 2018, Volume 11, Issue 6. 1923 <http://www.bioflux.com.ro/aac>

optimal. The

pH range of 6.8–7.4 in the ameliorant and lime treatments is still good enough for the growth of *C. striata* juveniles, because it is still in tolerance range of pH for *C. striata* life (4.25–9.4) (Courtenay & Williams 2004), while the optimal pH for the intensive fish farming is 7.0–8.0 (Wedemeyer 1996). In this pH range, gills can maximally bind oxygen and ions required by the body, so that the physiological activities of the fish will be more optimal. The use of ameliorant is related to the increase in Ca^{2+} needed to bind sulfate (SO_4^{2-}) (a source of the water acidity) to form gypsum deposits (Fitzpatrick et al 1998). Humic acid, fulvic acid and humin acid contained in an ameliorant cause an increase in pH (H^+ decreases). The sulfate level in ameliorant treatment was 10.2–12.6 mg L^{-1} , this range was still within the

the ability of the fish to absorb nutrient from the feed. It is reflected in the high albumin content in the fish reared in the ameliorant treated pond. By maintaining albumin in the blood plasma, the fish can also maintain blood volume. Albumin is one of blood plasma proteins synthesized in the liver (Infusino & Panteghini 2013). Albumin plays a role in transporting small molecules through plasma and extracellular fluid (Suprayitno 2014), maintaining plasma balance in blood vessels with fluid in the interstitial cavity and the formation of cells and tissues (Nugroho 2012).

AAFL Bioflux, 2018, Volume 11, Issue 6. 1924 <http://www.bioflux.com.ro/aac>

C. striata juveniles which were reared in the medium treated with an ameliorant had higher feed intake and feed efficiency than those of control, but those were not significantly different from the lime treatment. This showed that the feed given could be properly utilized by the fish, so that nutrients contained in the feed could be retained in the fish body efficiently. The feed efficiency is determined by several factors such as genetic, nutritional content, farming management, and environment (Robinson & Li 2015; Hasan & Soto 2017). In normal conditions, physiological processes in the fish's body will run well, so that the feed can be used entirely by the fish and can be converted into the fish flesh (Nelson & Chabot 2011). Daily feed intake is one of factors that influence the potential of the fish to grow optimally and feed consumption rate is closely related to the capacity and the emptying of the fish gut (Abowei & Ekubo 2011). The high SGR in the ameliorant treatment was not only

tolerance range for the fish life. In all treatments, the sulfate levels in the experimental mediums were low and still within the tolerance range for fish life according to Boyd (1988). The improvement of water quality in ponds located in acid sulfate land through the administration of an ameliorant produced from banana stems in *C. striata* farming provided the best results compared to the control and a treatment using lime, because the water's toxicity potential became reduced. This was shown by a lower ammonia (NH_3) level ($0.01\text{--}0.05 \text{ mg L}^{-1}$) in ameliorant treatment compared to other treatments. This level can still be tolerated by *C. striata*, because

at ammonia level of 1.57 mg L^{-1} , *C. striata* juvenile can still grow (Qin et al 1997).

Dissolved oxygen (DO) level range during the study was $4.12\text{--}5.80 \text{ mg L}^{-1}$. This range still meets the range recommended for the life and the growth of *C. striata* (Boyd 1988). *C. striata* is categorized in air breathing fish that has diverticula as the additional respiratory organ, so the fish can live in poor-oxygen-medium and can tolerate hypoxia by using oxygen derived from the air (Huang et al 2011; Song et al 2013). Alkalinity in water is caused by bicarbonate (HCO_3^-), carbonate (CO_3^{2-}), and hydroxide (OH^-) ions. At the ameliorant treatment, the alkalinity level was $85.1\text{--}115.8 \text{ mg L}^{-1} \text{ CaCO}_3$. This level was slightly higher than those of the lime treatment and control, and it was not far from the optimal range ($100\text{--}150 \text{ mg L}^{-1} \text{ CaCO}_3$) (Wedemeyer 1996). In the optimal range, the ability of alkalinity to sustain the pH of the water becomes better, but it does not

reflected in the albumin content and feed efficiency, but also reflected in protein retention (PR). The ameliorant and lime treatments resulted in higher PR (53.41% and 47.98%) than that of control (Table 2). Protein retention is affected by the type of the feed, feed quality and environmental conditions (Samantaray & Mohanty 1997; Purnamawati et al 2017a). The better is the feed quality and the environment, higher is the protein retention. The administration of an ameliorant and lime into the pond has a major role in improving water quality, so that the utilization of the feed energy becomes more optimal. This causes the protein derived from the feed will be retained more in the body and decomposed only in a few amount or it will be utilized as energy to maintain homeostasis. Jobling et al (1991) state that the high energy for body activities will reduce the energy budget for the growth. According to Nelson & Chabot (2011), energy in the feed is physiologically used for activities and metabolism, while the residue will be deposited as body tissue through the growth process and the synthesis of reproductive tissue. The difference in water quality conditions as a result of different treatments did not significantly influence fat retention ($p < 0.05$). This described that the utilization of fat by the fish in three experimental groups was not much different. This is thought to occur because the energy derived from the feed, especially derived from protein, is widely released by the body for the adaptation to poor environmental conditions. Conclusions. An ameliorant from banana stems can be used as a substitute for lime in improving water quality of a pond located in acid sulfate land. The use of an ameliorant and dolomite resulted in higher

negatively affect the fish growth (Wedemeyer 1996; Boyd 1988). The results of statistical analysis showed that the administration of an ameliorant and lime significantly affected the survival (SR) of *C. striata*. The administration of an ameliorant at a dose of 300 kg per ha pond and lime at a dose of 180 kg per ha pond produced a higher SR compared to control ($p > 0.05$). This was due to the better water quality in both treatment mediums compared to control. This caused the stress level of the fish decreasing and the fish became healthier. As a consequence, SR in ameliorant and lime treatments was also higher. Growth is a complex biological process. Growth can occur when there is excess energy and materials derived from the feed. The *C. striata* juveniles which were reared in the ameliorant treated pond produced the highest specific growth rate (SGR) (6.42%), followed by the lime and the control groups with SGR values of 5.54% and 4.77%, respectively ($p > 0.05$) (Table 2). The difference in growth rates showed that the *C. striata* which were reared in the ameliorant treated pond were better in utilizing the energy sourced by the feed. The high SGR in *C. striata* juveniles reared in the ameliorant treated pond was supported by a finding showing the highest albumin content in the fish blood (3.12 g per 100 mL) found in ameliorant treatment compared to other treatments ($p > 0.05$). A better water quality in the ameliorant treated pond will improve the ability of the fish to absorb nutrient from the feed. It is reflected in the high albumin content in the fish reared in the ameliorant treated pond. By maintaining albumin in the blood plasma, the fish can also maintain blood volume. Albumin is one of blood plasma proteins synthesized in

survivals of the *C. striata* than that of without using an ameliorant and dolomite, but the best growth was obtained in the ameliorant group. References Abowei J. F. N., Ekubo E. T., 2011 Some principles and requirements in fish nutrition. *British Journal of Pharmacology and Toxicology* 2:163–178. Alwi M., 2014 [The prospect of tidal swamp land for rice crops]. *The Proceeding of the National Seminar "Location-Specific Technological Innovation"*, 2014 August 6-7; Banjarbaru, Indonesia, pp. 45-59. [In Indonesian]. Arsyad D. M., Saidi B. B., Enrizal, 2014 [Development of agricultural innovations in tidal swamp land for increasing food sovereignty]. *Pengembangan Inovasi Pertanian* 7:169-178. [In Indonesian]. Boyd C. E., 1988 Water quality in warmwater fish ponds. Alabama Experiment Station, Auburn, 359 p. Cavalcante D. H., da Silva S. R., Pinheiro P. D., Akao M. M. F., e Sá M. V. C., 2012 Single or paired increase of total alkalinity and hardness of water for cultivation of Nile tilapia juveniles, *Oreochromis niloticus*. *Acta Scientiarum* 34:177–183. Courtenay Jr. W. R., Williams J. D., 2004 Snakeheads (Pisces, Channidae) - A biological synopsis and risk assesment. U.S. Geological Survey, Denver, 143 p.

AAFL Bioflux, 2018, Volume 11, Issue 6. 1925 <http://www.bioflux.com.ro/aafl> de Queiroz J. F., Nicolella G., Wood C. W., Boyd C. E., 2004 Lime application methods, water and bottom soil acidity in fresh water fish ponds. *Scientia Agricola* 61:469- 475. Fitzpatrick R. W., Merry R. H., Williams J., White I., Bowman G., Taylor G., 1998 Acid sulfate soil assessment: coastal, inland and minespoil conditions. National Land and Water

the liver (Infusino & Panteghini 2013). Albumin plays a role in transporting small molecules through plasma and extracellular fluid (Suprayitno 2014), maintaining plasma balance in blood vessels with fluid in the interstitial cavity and the formation of cells and tissues (Nugroho 2012).

AACL Bioflux, 2018, Volume 11, Issue 6. 1924 <http://www.bioflux.com.ro/aac>

C. striata juveniles which were reared in the medium treated with an ameliorant had higher feed intake and feed efficiency than those of control, but those were not significantly different from the lime treatment. This showed that the feed given could be properly utilized by the fish, so that nutrients contained in the feed could be retained in the fish body efficiently. The feed efficiency is determined by several factors such as genetic, nutritional content, farming management, and environment (Robinson & Li 2015; Hasan & Soto 2017). In normal conditions, physiological processes in the fish's body will run well, so that the feed can be used entirely by the fish and can be converted into the fish flesh (Nelson & Chabot 2011). Daily feed intake

is one of factors that influence the potential of the fish to grow optimally and feed consumption rate is closely related to the capacity and the emptying of the

fish gut (Abowei & Ekubo 2011). The high SGR in the ameliorant treatment was not only reflected in the albumin content and feed efficiency, but also reflected in protein retention (PR). The

Resources Audit Methods Paper. National Land and Water Resources Department, Canberra, 18 p. Gam L. H., Leow C. Y., Baie S., 2006 Proteomic analysis of snakehead fish (*Channa striata*) muscle tissue. *Malaysian Journal of Biochemistry and Molecular Biology* 14:25–32. Hasan M. R., Soto D., 2017 Improving feed conversion ratio and its impact on reducing greenhouse gas emissions in aquaculture. *FAO Non-Serial Publication, Food and Agriculture Organization of the United Nations, Rome*, 33 p. Hidayatullah S., Muslim, Taqwa F. H., 2015 [Rearing of snakehead larvae (*Channa striata*) in plastic lined pond with different stocking density]. *Jurnal Perikanan dan Kelautan* 20:61-70. [In Indonesian]. Huang C. Y., Lin C. P., Lin H. C., 2011 Morphological and biochemical variations in the gills of 12 aquatic air-breathing anabantoid fish. *Physiological and Biochemical Zoology* 84:125–134. Infusino I., Panteghini M., 2013 Serum albumin: accuracy and clinical use. *Clinica Chimica Acta* 419:15-18. Jobling M., Knudsen R., Pedersen P. S., dos Santos J., 1991 Effects of dietary composition and energy content on the nutritional energetics of cod, *Gadus morhua*. *Aquaculture* 92:243-257. Kang'ombe J., Brown J. A., 2008 Effect of salinity on growth, feed utilization, and survival of *Tilapia rendalli* under laboratory conditions. *Journal of Applied Aquaculture* 20:256-271. Marimuthu K., Jesu Arokiaraj A., Haniffa M. A., 2009 Effect of diet quality on seed production of the spotted snakehead *Channa punctatus* (Bloch). *American-Eurasian Journal of Sustainable Agriculture* 3:344–347. Mohd S. M. A., Abdul M. M. J., 2012 Therapeutic potential of the haruan (*Channa striatus*): from food to medicinal uses. *Malaysian Journal of Nutrition*

ameliorant and lime treatments resulted in higher PR (53.41% and 47.98%) than that of control (Table 2). Protein retention is affected by the type of the feed, feed quality and environmental conditions (Samantaray & Mohanty 1997; Purnamawati et al 2017a). The better is the feed quality and the environment, higher is the protein retention. The administration of an ameliorant and lime into the pond has a major role in improving water quality, so that the utilization of the feed energy becomes more optimal. This causes the protein derived from the feed will be retained more in the body

and decomposed only in a few amount or it will be utilized as energy to maintain homeostasis.

Jobling et al (1991) state that the high energy for body activities will reduce the energy budget for the growth. According to Nelson & Chabot (2011), energy in the feed is physiologically used for activities and metabolism,

while the residue will be deposited as body tissue through the growth process and the synthesis of reproductive tissue. The difference in water quality conditions as a result of different treatments did not significantly influence fat retention ($p < 0.05$). This described that the utilization of fat by the fish in three experimental groups was not much different. This is thought to occur because the energy derived from the feed, especially derived from protein, is widely released by the body for the adaptation to poor environmental conditions. Conclusions. An

18:125–136. Mollah M. F. A., 1985 Effects of stocking density and water depth on the growth and survival of freshwater catfish (*Clarias macrocephalus* Gunther) larvae. *Indian Journal of Fisheries* 35:1–17. Mollah M. F. A., Mamun M. S. A., Sarowar M. N., Roy A., 2009 Effects of stocking density on the growth and breeding performance of broodfish and larval growth and survival of shol, *Channa striatus* (Bloch). *Journal of the Bangladesh Agricultural University* 7:427-432. Nakkrasae L., Wisetdee K., Charoenphandhu N., 2015 Osmoregulatory adaptations of freshwater air-breathing snakehead fish (*Channa striata*) after exposure to brackish water. *Journal of Comparative Physiology B* 185:527–537. Nelson J. A., Chabot D., 2011 General energy metabolism. In: *Encyclopedia of fish physiology: from genome to environment*, volume 3. Farrell A. P. (ed), pp. 1566-1572, Academic Press, San Diego. Nugroho M., 2012 [The isolation of albumin and characteristics of molecular weight from extraction result by steaming of fish gabus (*Ophiocephalus striatus*)]. *SAINTEK PERIKANAN: Indonesian Journal of Fisheries Science and Technology* 9:40–48. [In Indonesian]. Purnamawati, 2017 [The growth performance of snakehead *Channa striata* Bloch. on tidal land through water quality engineering]. Bogor Agricultural University, Bogor, 78 p. [In Indonesian]. Purnamawati, Djokosetiyanto D., Nirmala K., Harris E., Affandi R., 2017a Survival and growth of striped snakehead fish (*Channa striata* Bloch.) juvenile reared in acid sulfate water and rainwater medium. *AACL Bioflux* 10:265–273. Purnamawati, Djokosetiyanto D., Nirmala K., Surawidjaja E. H., Affandi R., 2017b Survival and

ameliorant from banana stems can be used as a substitute for lime in improving water quality of a pond located in acid sulfate land. The use of an ameliorant and dolomite resulted in higher survivals of the *C. striata* than that of without using an ameliorant and dolomite, but the best growth was obtained in the ameliorant group. References

Abowei J. F. N., Ekubo E. T., 2011 Some principles and requirements in fish nutrition. *British Journal of Pharmacology and Toxicology* 2:163–178.

Alwi M., 2014 [The prospect of tidal swamp land for rice crops]. The Proceeding of the National Seminar "Location-Specific Technological Innovation", 2014 August 6-7; Banjarbaru, Indonesia, pp. 45-59. [In Indonesian]. Arsyad D. M., Saidi B. B., Enrizal, 2014 [Development of agricultural innovations in tidal swamp land for increasing food sovereignty]. *Pengembangan Inovasi Pertanian* 7:169-178. [In Indonesian].

Boyd C. E., 1988 Water quality in warmwater fish ponds. Alabama Experiment Station, Auburn, 359

p. Cavalcante D. H., da Silva S. R., Pinheiro P. D., Akao M. M. F., e Sá M. V. C., 2012 Single or paired increase of total alkalinity and hardness of water for cultivation of Nile tilapia juveniles, *Oreochromis niloticus*. *Acta Scientiarum* 34:177–183.

Courtenay Jr.

growth response of snakehead fish *Channa striata* Bloch juvenile in

AAFL Bioflux, 2018, Volume 11, Issue 6. 1926 <http://www.bioflux.com.ro/aafl> aerated and unaerated acid sulfate water. *Jurnal Akuakultur Indonesia* 16:60–67. Qin J., Fast A. W., Kai A. T., 1997 Tolerance of snakehead *Channa striatus* to ammonia at different pH. *Journal of the World Aquaculture Society* 28:87-90. Rahman M. A., Arshad A., Amin S. M. N., Shamsudin M. N., 2013 Growth and survival of fingerlings of a threatened snakehead, *Channa striatus* (Bloch) in earthen nursery ponds. *Asian Journal of Animal and Veterinary Advances* 8:216–226. Robinson E. H., Li M. H., 2015 Feed conversion ratio for pond-raised catfish. Information Sheet No. 1364. The Office of Agricultural Communications, The Division of Agriculture, Forestry, and Veterinary Medicine, Mississippi State University, Strakville, 4 p. Samantaray K., Mohanty S. S., 1997 Interactions of dietary levels of protein and energy on fingerling snakehead, *Channa striata*. *Aquaculture* 156:241–249. Song L. M., Munian K., Rashid Z. A., Bhassu S., 2013 Characterization of Asian snakehead murrel *Channa striata* (Channidae) in Malaysia: an insight into molecular data and morphological approach. *The Scientific World Journal* 2013, Article ID 917506, <http://dx.doi.org/10.1155/2013/917506>. Suprayitno E., 2014 Profile albumin fish corks (*Ophicephalus striatus*) of different ecosystems. *International Journal of Current Research and Academic Review* 2:201–208. Takeuchi T., 1988 Laboratory work-chemical evaluation of dietary nutrient. In: *Fish nutrition and*

W. R., Williams J. D., 2004 Snakeheads (Pisces, Channidae) - A biological synopsis and risk assesment. U.S. Geological Survey, Denver, 143 p.

AAFL Bioflux, 2018, Volume 11, Issue 6. 1925 <http://www.bioflux.com.ro/aacld>

de

Queiroz J. F., Nicolella G., Wood C. W., Boyd C. E., 2004 Lime application methods, water and bottom soil acidity in fresh water fish ponds. *Scientia Agricola* 61:469- 475. Fitzpatrick R. W., Merry R. H., Williams J., White I., Bowman G., Taylor G., 1998 Acid sulfate soil assessment: coastal, inland and minespoil conditions. National Land and Water Resources Audit Methods Paper. National Land and Water Resources Department, Canberra, 18 p. Gam L. H., Leow C. Y., Baie S., 2006

Proteomic analysis of snakehead fish (*Channa striata*) muscle tissue. *Malaysian Journal of Biochemistry and Molecular Biology* 14:25–32.

Hasan M. R., Soto D., 2017 Improving feed conversion ratio and its impact on reducing greenhouse gas emissions in aquaculture. FAO Non-Serial Publication, Food and Agriculture Organization of the United Nations, Rome, 33 p.

Hidayatullah S., Muslim, Taqwa F. H., 2015 [Rearing of snakehead larvae (*Channa striata*) in plastic lined pond with different

mariculture. Watanabe T. (ed), pp. 179-233, Kanagawa International Fisheries Training Centre, Japan International Cooperation Agency (JICA), Tokyo. Vivekanandan E., 1977 Surfacing activity and food utilization in the obligatory air-breathing fish *Ophiocephalus striatus* as a function of body weight. *Hydrobiologia* 55:99-112. Weatherley A. H., Gill H. S., 1989 The biology of fish growth. Academic Press, London, 443 p. Wedemeyer G. A., 1996 Physiology of fish in intensive culture system. Springer US, New York, 232 p. Wedemeyer G. A., Yasutake W. T., 1977 Clinical methods for the assessment of the effects of environmental stress on fish health. U.S. Fish and Wildlife Service, Washington D.C., 18 p. *** APHA (American Public Health Association), 1989 Standard methods for the examination of water and wastewater 17 th edition. American Public Health Association, Washington D.C., 1268 p. *** BPS Kalbar (Badan Pusat Statistik Provinsi Kalimantan Barat), 2012 [West Kalimantan in number]. Badan Pusat Statistik Provinsi Kalimantan Barat, Pontianak. [In Indonesian]. Received: 01 November 2018. Accepted: 02 December 2018. Published online: 10 December 2018. Authors: Purnamawati, Pontianak State Polytechnic, Department of Marine Science and Fisheries, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: pur_polnep@yahoo.com Nurmala, Pontianak State Polytechnic, Department of Marine Science and Fisheries, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: ibu.nurmala@gmail.com Mochammad Idham Shilman, Pontianak State Polytechnic, Department of Marine Science and

stocking density]. *Jurnal Perikanan dan Kelautan* 20:61-70. [In Indonesian].

Huang C. Y., Lin C. P., Lin H. C., 2011 Morphological and biochemical variations in the gills of 12 aquatic air-breathing anabantoid fish. *Physiological and Biochemical Zoology* 84:125-134.

Infusino I., Panteghini M., 2013 Serum albumin: accuracy and clinical use. *Clinica Chimica Acta* 419:15-18.

Jobling M., Knudsen R., Pedersen P. S., dos Santos J., 1991 Effects of dietary composition and energy content on the nutritional energetics of cod, *Gadus morhua*. *Aquaculture* 92:243-257.

Kang'ombe J., Brown J. A., 2008 Effect of salinity on growth, feed utilization, and survival of *Tilapia rendalli* under laboratory conditions. *Journal of Applied Aquaculture* 20:256-271.

Marimuthu K., Jesu Arokiaraj A., Haniffa M. A., 2009 Effect of diet quality on seed production of the spotted snakehead *Channa punctatus* (Bloch). *American-Eurasian Journal of Sustainable Agriculture* 3:344-347.

Mohd S. M. A., Abdul M. M. J., 2012

Therapeutic potential of the haruan (*Channa striatus*): from food to medicinal uses.

Malaysian Journal of Nutrition 18:125-136.

Fisheries, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: idham.shilman@yahoo.co.id Eko Dewantoro, Muhammadiyah University of Pontianak, Faculty of Fisheries and Marine Science, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: ekodewe_ump@yahoo.com Diah Ayu Satyari Utami, Marine and Fisheries Polytechnic of Jembrana, Study Program of Aquaculture, Indonesia, Bali, Jembrana 82218, Desa Pengambangan, e-mail: dplongitm@gmail.com This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited. How to cite this article: Purnamawati, Nurmala, Shilman M. I., Dewantoro E., Utami D. A. S., 2018 Effects of ameliorant on the growth of snakehead fish (*Channa striata*) juvenile reared in acid sulfate water medium located in tidal land. *AAFL Bioflux* 11(6):1919-1926.

Mollah M. F. A., 1985 Effects of stocking density and water depth on the growth and survival of freshwater catfish (*Clarias macrocephalus* Gunther) larvae. *Indian Journal of Fisheries* 35:1-17.

Mollah M. F. A., Mamun M. S. A., Sarowar M. N., Roy A., 2009 Effects of stocking density on the growth and breeding performance of broodfish and larval growth and survival of shol, *Channa striatus* (Bloch). *Journal of the Bangladesh Agricultural University* 7:427-432.

Nakkrasae L., Wisetdee K., Charoenphandhu N., 2015

Osmoregulatory adaptations of freshwater air-breathing snakehead fish (*Channa striata*) after exposure to brackish water.

Journal of Comparative Physiology B 185:527-537.

Nelson J. A., Chabot D., 2011 General energy metabolism. In: *Encyclopedia of fish physiology: from genome to environment*, volume 3. Farrell A. P. (ed), pp. 1566- 1572, Academic Press, San Diego.

Nugroho M., 2012 [The isolation of albumin and characteristics of molecular weight from extraction result by steaming of fish gabus (*Ophiocephalus striatus*)]. *SAINTEK PERIKANAN: Indonesian Journal of Fisheries Science and Technology* 9:40-48. [In Indonesian].

Purnamawati, 2017 [The growth performance of snakehead *Channa striata* Bloch. on tidal land through water quality engineering]. Bogor Agricultural University, Bogor, 78 p. [In Indonesian].

Purnamawati, Djokosetiyanto D., Nirmala K., Harris E., Affandi R., 2017a

Survival and growth of striped snakehead fish (*Channa striata* Bloch.) juvenile reared in acid sulfate water and rainwater medium. *AAFL Bioflux* 10:265–273.

Purnamawati, Djokosetiyanto D., Nirmala K., Surawidjaja E. H., Affandi R., 2017b

Survival and growth response of snakehead fish *Channa striata* Bloch juvenile

in

AAFL Bioflux, 2018, Volume 11, Issue 6. 1926 <http://www.bioflux.com.ro/aal>

aerated and unaerated acid sulfate water.

Jurnal Akuakultur Indonesia 16:60–67.

Qin J.,

Fast A. W., Kai A. T., 1997 Tolerance of snakehead *Channa striatus* to ammonia at different pH. *Journal of the World Aquaculture Society* 28:87-90.

Rahman M. A., Arshad A.,

Amin S. M. N., Shamsudin M. N., 2013 Growth and survival of fingerlings of a threatened snakehead, *Channa striatus* (Bloch) in earthen nursery ponds. *Asian Journal of Animal and Veterinary Advances* 8:216–226. Robinson E. H., Li M. H., 2015 Feed conversion ratio for pond-raised catfish. Information Sheet No. 1364. The Office of Agricultural Communications, The

Division of Agriculture, Forestry, and Veterinary Medicine, Mississippi State University,

Strakville, 4 p.

Samantaray K., Mohanty S. S., 1997

Interactions of dietary levels of protein and energy on fingerling snakehead, *Channa striata*. *Aquaculture* 156:241–249.

Song L. M., Munian K., Rashid Z. A., Bhassu S., 2013
Characterization

of Asian snakehead murrel *Channa striata* (Channidae) in Malaysia: an insight into molecular data and morphological approach.

The

Scientific World Journal 2013, Article ID 917506, <http://dx.doi.org/10.1155/2013/917506>.

Suprayitno E., 2014 Profile albumin fish cork (*Ophicephalus striatus*) of different ecosystems. *International Journal of Current Research and Academic Review* 2:201– 208.

Takeuchi T., 1988 Laboratory work-chemical evaluation of dietary nutrient. In: *Fish nutrition and mariculture*. Watanabe T. (ed), pp. 179-233, Kanagawa International Fisheries Training Centre, Japan International Cooperation Agency (JICA), Tokyo.

Vivekanandan E., 1977 Surfacing activity and food utilization in the obligatory air-breathing fish *Ophiocephalus striatus* as a function of body weight. *Hydrobiologia* 55:99-112.

Weatherley A. H., Gill H. S., 1989 *The biology of fish growth*. Academic Press, London, 443

p. Wedemeyer G. A., 1996 *Physiology of fish in intensive culture system*. Springer US, New York, 232 p.

Wedemeyer G. A., Yasutake

W. T., 1977

Clinical methods for the assessment of

the effects of environmental stress on fish health.

U.S. Fish

and Wildlife Service,

Washington D.C., 18

p. ***

APHA (

American Public Health Association), 1989

Standard methods for the examination of water and wastewater
17 th

edition.

American Public Health Association,

Washington D.C., 1268

p. *** BPS Kalbar (Badan Pusat Statistik Provinsi Kalimantan Barat), 2012 [West Kalimantan in number]. Badan Pusat Statistik Provinsi Kalimantan Barat, Pontianak. [In Indonesian]. Received: 01 November 2018. Accepted: 02 December 2018. Published online: 10 December 2018. Authors: Purnamawati,

Pontianak State Polytechnic, Department of Marine Science and Fisheries, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: pur_polnep@yahoo.com Nurmala, Pontianak State Polytechnic,

Department of Marine Science and Fisheries, Study Program of Aquaculture, Indonesia, West

Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: ibu.nurmala@gmail.com Mochammad Idham Shilman,

Pontianak State Polytechnic, Department of Marine Science and Fisheries, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail:

idham.shilman@yahoo.co.id Eko Dewantoro, Muhammadiyah University of Pontianak, Faculty of Fisheries and

Marine Science, Study Program of Aquaculture, Indonesia, West Kalimantan, Pontianak 78124, Jalan Ahmad Yani, e-mail: ekodewe_ump@yahoo.com

Diah Ayu Satyari Utami, Marine and Fisheries Polytechnic of Jembrana, Study Program of Aquaculture, Indonesia, Bali, Jembrana 82218, Desa Pengambangan,

e-mail: dplongitm@gmail.com

This is an open-access article distributed under the terms of

the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original

author and source are credited. How to cite this article:

Purnamawati,

Nurmala, Shilman M. I., Dewantoro E., Utami D. A. S., 2018

Effects of ameliorant on the

growth of snakehead fish (*Channa striata*) juvenile reared in acid sulfate water

medium located in tidal land. AACL Bioflux 11(6):1919-1926.
